

[ZP2008(11-10)]

November 10, 2008

*" Never shoot, Never hit "***THE CANCELLATION PROBLEM FOR AFFINE SPACES**

SUSUMU ODA

ABSTRACT. The Cancellation Problem for Affine Spaces is settled affirmatively, that is, it is proved that : Let k be an algebraically closed field of characteristic zero and let $n, m \in \mathbb{N}$. If $R[Y_1, \dots, Y_m] \cong_k k[X_1, \dots, X_{n+m}]$ as k -algebras, where $Y_1, \dots, Y_m, X_1, \dots, X_{n+m}$ are indeterminates, then $R \cong_k k[X_1, \dots, X_n]$.

The Cancellation Problem for Affine Spaces (or Zariski Problem) is the following:

The Cancellation Problem for Affine Spaces(or Zariski Problem). *Let k be an algebraically closed field of characteristic zero and let $n, m \in \mathbb{N}$. If $R[Y_1, \dots, Y_m] \cong_k k[X_1, \dots, X_{n+m}]$ as k -algebras, where $Y_1, \dots, Y_m, X_1, \dots, X_{n+m}$ are indeterminates, is $R \cong_k k[X_1, \dots, X_n]$?*

We may assume that $k = \mathbb{C}$. in the above problem. The answer is yes for $n = 1$ ([8]) or $n = 2$ ([4]). But the cases $n \geq 3$ are open problems. Moreover, the known case $n = 1, 2$ are proved by some tools in algebraic geometry. The purely algebraic proof has not been known yet even in such cases.

Our objective is to give an affirmative algebraic proof for all $n \in \mathbb{N}$.

Throughout this paper, all fields, rings and algebras are assumed to be commutative with unity. For a ring R , $R^\times$ denotes the set of units of R and $K(R)$ the total quotient ring. $\text{Spec}(R)$ denotes the affine scheme defined by R or merely

2000 Mathematics Subject Classification : Primary 13C25, Secondary 15A18

Key words and phrases: The Cancellation Problem for Affine Spaces, unramified, etale, polynomial rings and affine spaces

the set of all prime ideals of R and $\text{Ht}_1(R)$ denotes the set of all prime ideals of height one. Our general reference for unexplained technical terms is [6].

1. PRELIMINARIES

Definition 1.1. Let $f : A \rightarrow B$ be a ring-homomorphism of finite type of locally Noetherian rings. The homomorphism f is called *unramified* if $PB_P = (P \cap A)B_P$ and $k(P) = B_P/PB_P$ is a finite separable field extension of $k(P \cap P) = A_{P \cap P}/(P \cap A)A_{P \cap P}$ for all prime ideal P of B . The homomorphism f is called *etale* if f is unramified and flat.

Remark 1.2. In the above Definition 1.1, we have only to consider maximal ideals M of A instead of prime ideals P of A (see, [1]).

Proposition 1.3. ([10]) *Let k be an algebraically closed field of characteristic zero and let B be a polynomial ring $k[Y_1, \dots, Y_n]$. Let D be an extension of B . If D is etale and finite over B then $D = B$.*

The proof of the following is seen in a lot of texts of algebra, e.g., [9].

Lemma 1.4. *Let k be an algebraically closed field of characteristic zero and let $f(X)$ be a non-constant polynomial in $k[X_1, \dots, X_n][X]$. Then there exist $c_1, \dots, c_n \in k$ such that, putting $X'_i = X_i + c_iX$ ($1 \leq i \leq n$), $f(X)$ is a monic polynomial in $k[X'_1, \dots, X'_n][X]$.*

Let A be a Krull domain and let $Cl(A)$ denote the divisor class group of A .

Lemma 1.5. ([2, Theorem 8.1]) *Let R be a Krull domain and let $R[X]$ be a polynomial ring. Then $Cl(R) \cong Cl(R[X])$.*

Lemma 1.6. ([2, Proposition 6.1]) *Let R be a Krull domain. Then R is a UFD if and only if $Cl(R) = 0$.*

Corollary 1.7. *If R be a Krull domain and let $R[X]$ be a polynomial ring. If $R[X]$ is a UFD, then R is a UFD.*

Lemma 1.8. ([7, Theorem 14.2]) *Let (R, m) be an n -dimensional regular local ring, and let $x_1, \dots, x_i$ be elements of m . Then the following conditions are equivalent :*

- (1) $x_1, \dots, x_i$ is a subset of a regular system of parameters of R ;
- (2) the images in m/m^2 of $x_1, \dots, x_i$ are linear independent over a field R/m ;
- (3) $R/(x_1, \dots, x_i)$ is an $(n - i)$ -dimensional regular local ring.

Lemma 1.9. *If R is a k -affine domain over an algebraically field k and if $R[X]$ is a polynomial ring over k , then R is a regular domain, that is, R is regular $\iff R[X]$ is regular.*

Proof. Since any maximal ideal of $R[X]$ is a form $mR[X] + (X - a)R[X]$ for some $a \in k$ and some maximal ideal m of R , mR_m has a regular system of parameters. Hence R_m is regular for all maximal ideal m of R . Thus R is regular. $\square$

Proposition 1.10. (c.f. [7, Theorem 25.2]) *Let k be an algebraically closed field of characteristic zero, let R be a regular k -affine domain and let $R[X]$ be a polynomial ring and let $f(X) \in R[X] \setminus R$ be a monic polynomial in X . Assume that $f(X)$ is a member of a regular system of parameters of $MR[X]_M$ such that $f(X) \in M \setminus (M^2 + (m \cap R)R[X])$ for each maximal ideal M of $R[X]$ such that $f(X) \in M$*

(since R is regular, $R[X]$ is also regular by lemma 1.9). Then $R[X]/f(X)R[X]$ is unramified over R .

Proof. Let M be a maximal ideal of $R[X]$ which satisfies the assumption in Proposition. Put $m = M \cap R$, which is a maximal ideal of R because R is a k -affine domain. Then $\bar{f} \in R/m[X] = k[X]$ is a regular system of parameters of $k[X]_{\bar{M}}$, where $\bar{M}$ is the image of M in $R/m[X]$. Hence $\bar{M}k[X]_{\bar{M}} = \bar{f}k[X]_{\bar{M}}$ by the property of $f(X)$. Thus $m(R[X]_M/fR[X]_M) = (M/fR[X]_M)(R[X]/fR[X]_M)$. Since M is any maximal ideal of $R[X]$ containing $f(X)$, we conclude that $R[X]/f(X)R[X]$ is unramified over R . $\square$

2. MAIN RESULT

Now we prove our main result, the Cancellation Problem for Affine Spaces.

Theorem 2.1 (The Cancellation Problem for Affine Spaces). *Let k be an algebraically closed field of characteristic zero and let $n, m \in \mathbb{N}$. If $R[Y_1, \dots, Y_m] \cong_k k[X_1, \dots, X_{n+m}]$ as k -algebras, where $Y_1, \dots, Y_m, X_1, \dots, X_{n+m}$ are indeterminates, then $R \cong_k k[X_1, \dots, X_n]$.*

In order to prove this theorem, we may assume that $m = 1$ by induction. So we have only to show the following :

Theorem 2.2 (Special Case). *If $R[T] \cong_k k[X_1, \dots, X_n, X_{n+1}]$, then $R \cong_k k[X_1, \dots, X_n]$.*

Proof. Put $\phi : R[T] \cong_k k[X_1, \dots, X_{n+1}]$. First note that R is a regular UFD by Corollary 1.7 and Lemma 1.9. Put $\phi(T) = f(X_1, \dots, X_{n+1}) \in k[X_1, \dots, X_{n+1}]$, which is irreducible in $k[X_1, \dots, X_{n+1}]$. Furthermore $f(X_1, \dots, X_{n+1}) \in k[X_1, \dots, X_{n+1}]$ is monic in X_{n+1} with coefficients in $k[X_1, \dots, X_n]$.

We can write $\phi(T) = f(X_1, \dots, X_{n+1}) \in k[X_1, \dots, X_n][X_{n+1}]$. We may also assume that $f(X_1, \dots, X_n X_{n+1}) \in k[X_1, \dots, X_n, X_{n+1}]$ in X_{n+1} by Lemma 1.4. Then

$$R \cong_k k[X_1, \dots, X_{n+1}]/(f(X_1, \dots, X_{n+1})) = k[x_1, \dots, x_n][\alpha],$$

where x_i denotes the residue class of X_i modulo $(f(X_1, \dots, X_{n+1}))$ and α is that of X_{n+1} . It is easy to see that $x_1, \dots, x_n$ are algebraically independent over k . Put $A := k[x_1, \dots, x_n]$, which is a polynomial ring. We may assume that $A \hookrightarrow D$ by the isomorphism $R \cong A[\alpha](\supseteq A)$. Note that any maximal ideal M' of A is given by $M \cap A$ for a maximal ideal M of R .

Note that $f(x_1, \dots, x_n, X) =: f(X) \in A[X]$ is irreducible in $A[X]$ and $f(\alpha) = 0$ and that $A[\alpha] = A[X]/(f(X)) \cong_k k[X_1, \dots, X_{n+1}]/(f(X_1, \dots, X_{n+1})) \cong_k R$, which is a regular domain by Lemma 1.9.

Since for any maximal ideal N of $R[T]$ containing T , $R[T]_N/TR[T]_N \cong R_{N \cap R}$ is regular, T is a member of a regular system of parameters of $NR[T]_N$ by Lemma 1.8. Since $T \in (N^2 + (N \cap R)R[T])$, it follows that $f(X) \in N'A[X]_{N'}$ satisfies $f(X) \in N' \setminus (N'^2 + (N' \cap A)A[X])$ for any maximal ideal N' of $A[X]$ containing $f(X)$. Hence $A[\alpha] \cong_A A[X]/f(X)A[X]$ is unramified over A by Proposition 1.10. Since $f(X) \in A[X]$ is monic, $A[X]/f(X)A[X]$ is flat over A . So $A[\alpha]$ is finite etale over A . Hence we have $A = A[\alpha] \cong_k R$ by Proposition 1.3. Therefore $R \cong_k A[\alpha] = A \cong_k k[X_1, \dots, X_n]$. $\square$

REFERENCES

- [1] A.Altman and S.Kleiman, Introduction to Grothendieck duality theory, Lecture Notes in Math. 146, Springer-Verlag (1970).
- [2] R.M.Fossum, The Divisor Class Group of a Krull Domain, Springer-Verlag(1973).
- [3] H.Bass, E.H.Connel and D.Wright, The Jacobian conjecture, Bull. A.M.S., 7(2) (1982),287-330.
- [4] T.Fujita, On Zariski Problem, Pro. Japan Acad. Ser. A Math. Sci 55 (1975), 106-110.
- [5] S.Lang, Algebra, Addison-Wesley(Sixth Printing) (1974).
- [6] H.Matsumura, Commutative Algebra, Benjamin, New York (1970).
- [7] H.Matsumura, Commutative Ring Theory, Cambridge Univ. Press(1980).
- [8] M.Miyanishi and T.Sugie, Affine surfaces containing cylinderlike open set, J. Math. Kyoto Univ. 20 (1980), 1-42.
- [9] M.Nagata, Local Rings, Interscience (1962).

- [10] M.Raynaud, Anneaux Locaux Henséliens, Lecture Notes in Math. 169, Springer-Verlag (1970).
- [11] D.Wright, On the Jacobian conjecture, Illinois J. Math., 25 (1981), 423-440.

Department of Mathematics

Faculty of Education

Kochi University

2-5-1 Akebono-cho, Kochi 780-8520

JAPAN

ssmoda@kochi-u.ac.jp